


FIG. 1

2/15



Total Residues.....55  
Basic Residues.....15  
Acid Residues.....4  
Net Charge.....+11  
IEP.....10.77

Molecular  
Weight.....6451.5  
  
280nm Molar  
Absorption.....6050



**FIG. 2**

3/15





FIG. 3




5/15



Reaction conditions: 52 mM p7NC + 744 mM NEM; pH 7.0, 60min. at RT.  
The positions of alkylated Cys residues were determined by sequence analysis of separated proteins and are indicated by the notation C#-M.

**FIG. 5**

6/15



The reactive functional groups are shaded

**FIG. 6**

7/15


disulfides




nitroso compounds




maleimides



$\alpha$ -halogenated ketones



phenylhydrazids



Nitric Oxide and Derivatives NO


cupric ions and complexes  $Cu^{+2}$

ferric ions and complexes  $Fe^{+3}$

where R is any atom or molecule, and X is selected from the group consisting of F, I, Br and Cl.

**FIG. 7**

8/15



**FIG. 8**



9/15






FIG. 9

10/15




**FIG. 10**

11/15



**FIG. 11**



3 Tetraethylthiuram Disulfide +  $\text{p7NC}$   $\rightarrow$  Oxidized p7 (3 S-S) + 6 Diethylthiocarbamate +  $2 \text{Zn}^{+2}$



4 Diethylthiocarbamate +  $2 \text{Zn}^{+2} \rightarrow 2$  Coordination Complexes

13/15



**Control**  
buffer + p7NC



**Imuthiol**  
(Diethylthiocarbamate)





**Disulfiram**  
(Tetraethylthiuram Disulfide)





**FIG. 13**

14/15


## Synthesis



## General Reactions



**FIG. 14**



15/15




FIG. 15